

;
 * ;
 * ;
 120 -
 43,9 86,6%
 41-50 ,
 35% > 40 ,
 0,8% (1).
 6,7%
 ST,
 - 120
 43,9 86,6%
 41-50 ,
 35% > 40 ,
 0,8% (1). 6,7%
 ST,
 () -
 - ().
 ST
 ().
 [1].
 [2]. - () ()
 ().
 -
 120 -
 20 , - 21-30 .. - 31-40 .. - 41-50 .. - 51-60 .
 - 60 . , 75%
 ()

20-29 . - 161/ , 30-39 . - 156/ , 40-49 . - 152/ , 50-69 . - 145/ [3].
 25
 (150 / 1) 25 3
 9-12
 60 1 .
 - , . , . ,
 , -02 (. , « »),
 6- (-).
 - , 1. -43,9 .
 I - -

20 .	5 (4,2)	5 (100%)	-	-	-	-5
21-30 .	19 (16)	18 (94,7)	-	1 (5,3)	-	-18, -1
31-40 .	41 (34)	36 (87,8)	-	-	5 (12,2)	-40, -1
41-50 .	35 (29)	30 (85,7)	-	2 (5,7)	3 (8,2)	-30, -2, -3
51-60 .	19 (16)	15 (78,9)	-	4 (21,1)	-	-15, -4
> 60 .	1 (08)	-	-	1 (100)	-	-1
:	120 (100)	104 (86,6)	-	8 (6,7)	8 (6,7)	

31-50 ., 63%
 20 21-30 . ,
 , 23 24 (95,8%)
 ().
 22- 125 136/ . ,
 ().
 31-40 .
 36 (87,8%)
 37 , Q-
 (). 150 158/
 5 (12,2%).
 150 150/
 156/ V₆ ST 2

150 ()
 156/
 ST. ()
 36- ,
 50 « »
 (). 2
 35 41-50 ,, , 30 (85,7%)
 75 143/ , ST 152/ 3- V₄₋₆
 75 94/ 15
 100 158/ , 2
 , 8,2%
 , 1 (0,8%) 42 .
 80/50
 51-60 ,, 19
 , 15 (78,9%) 25 125/ ()
 4 145/) ST 4
 V₅₋₆ 212. 6-
 3 25 1 10
 ST V₄₋₆ 2-3 .
 1 3 125 136/
 ST 1,5 .
 , 2-
 , 4- , 75
 - 150/ .
 10
 -
 60 1
 125/ (145/), 100
 ST 2,5 1-

3

1.

0,8%.

2.

41-50

- 35%

3.

(86,6%)

- 75%

- 85-90% (Chuhg E.X., 1979),

« »

4.

SUMMARY

**BICYCLE TESTING AND ITS ESTIMATION ABILITY OF THE
FUNCTIONAL STATE OF THE PATIENTS WITH CARDIAC AND NON- CARDIAC DISEASES.
SIGNIFICANCE OF THE HEART RATE AND BLOOD PRESSURE CHANGE**

V.V. Laba, O.V. Laba*

Sumy State University, Medical Institute, Sumy;

**Kharkiv Medical Academy of Postgraduate Education, Kharkiv*

Diagnostic possibilities bicycleergometer were studied in different age categories and diagnostic groups of 120 male patients. The average age of surveyed comprised 43,9. True negative result was achieved in 86,6% of patients. Since the age group 41-50 years increased the probability true positive test result, wich amounted to 35% in patients older than 40 years. The percentage of transient complications of hypotensive reaction renewable period was 0,8% (in 1 patient). Mistaken positive results were 6,7%.

Key words: bicycleergometer, stress test, ischemic heart disease, segment ST, heard rate.

1.

, 2006. - 90 .

2.

, 2002. - 296 .

3.

, 1990. - 192 .

12 2010 .