

BRONCHIAL ASTHMA AND OBESITY ASSOCIATION: PATHOGENETIC MECHANISM OF HARMFUL INFLUENCE REALIZATION AND QUALITY OF LIFE

V. V. Kmyta, *NR-1/1*

A. M. Dyadechko, *ELA*

The goal of the research was the comparative study of the level of inflammation markers during the isolated cases of bronchial asthma (BA) and when it was accompanied with visceral obesity (VO) and quality of life (LQ)

Materials and methods of investigation. The diagnostics of VO was conducted according to the recommendations of World Health Organization (WHO). The investigation of inflammation markers was conducted for 71 patients with BA, among which: Group 1 included 35 BA patients with normal body weight, and Group 2 had 36 VO patients. The Control Group consisted of 22 practically healthy people. The level of interleukins (IL) IL-6, IL-8 and the tumor necrosis factors- α (TNF- α) in blood serum was determined using immunofluorescence (IFH) method, the antigenic composition of the membrane of mononuclears in peripheral blood was determined using indirect immunofluorescent method, the level of α_1 -inhibitor proteinase (α_1 -IP) and α_2 -macroglobulin (α_2 -MG) was determined using spectrophotometric method. Quality of life was measured with the help of Asthma Quality of Life Questionnaire (AQLQ). The achieved results were analyzed using the methods of variance statistics. The correctness of the difference in the average measurements was evaluated using Student's t - criterion.

Results of investigation. Group 1 patients had 3x and Group 2 patients 8x the level of FNP- α , and IL-6 level was elevated by 2,5 and 5,6, respectively. The level of IL-6 in blood serum was abnormal for both patient groups compared to the Controls, while its level for VO patients was evidently higher compared to Group 1 patients ($p < 0,05$).

Group 2 patients had levels of CD3, CD4, CD8 and CD16 evidently lower compared to Group 1 patients ($p < 0,05$). The levels of adhesive molecules (CD11b, CD54), expression of receptor to IgE (CD23), activation markers of lymphocytes (receptors to IL-2, HLA-DR) were evidently higher for patients of both groups compared to the Control, and higher for Group 2 patients compared to Group 1. It was observed that α_1 -IP and α_2 -MG were elevated by 2x and 2,6x respectively compared to healthy individuals.

Results of investigation convey that the presence of VO in BA patients leads to: hyperproduction of proinflammation cytokines (IL-6, IL-8, TNF- α); increase of T - lymphopenia, deficiency of cytotoxic cells, increased expression of activation markers, adhesive molecules, receptors to IgE, HLA-DR; increased level of α_1 -IP and α_2 -MG proteins.

Study of LQ showed that in patients with obesity the level of physical activity ($p < 0.05$) and emotional status ($p < 0.05$) were lower along with a more pronounced influence of the environment ($p < 0.05$). Overall rating of LQ were in patients of I group (4.5 ± 0.4) points against (3.4 ± 0.3) points in II group ($p < 0.05$).

Conclusions. These changes, along with known negative impacts of obesity on the process of BA (bronchial hyperactivity, increased frequency of gastroesophageal reflux, aggravation of pulmonary deficiency), represent one more pathogenetic mechanism – the exacerbation of inflammatory process. Presence of obesity complicated BA duration, entailing a more pronounced reduction LQ that dictates new conditions of including in their process. We can make conclusions that level of inflammation markers is higher in group of patients with bronchial asthma and obesity, as a obesity tissues is an active endocrine organ which produce different types of inflammatory factors including proinflammatory cytokines.

Соціально-гуманітарні аспекти розвитку сучасного суспільства : матеріали IV Всеукраїнської наукової конференції викладачів, аспірантів, співробітників та студентів факультету іноземної філології та соціальних комунікацій, м. Суми, 19-20 квітня 2013 р. / Відп. за вип. В.В. Опанасюк. — Суми : СумДУ, 2013. — Ч.4. — С. 49-50.