```
// Azerbaijan medical
journal. -2014. - 1. - .92-95.
              50
           (n=26)
           (n=24)
                                             50
                                                 [1].
                              23%
           59%
                                        [2].
          [3].
```

```
( - ( ( -1 ),
                                           - 6 ( -6)) [4],
),
                     [5].
                                               [6].
                          [7].
                    -6,
           [8],
                                                         [9].
   -1
                                            [10].
                                                 )/
                                     50
                                                ( CR, 1990, 1991),
                             (1999).
                                                     ,
150
                                                                     12
-14
 (n=26) -
                                 (500
1
                                        /500
                                                              (n=24) -
                              20
```

: 3,0 (<0,001) 1.7 , (₋ <0,001).

3 n=20n=26 n=26 n=24n=2471,6±3,6* 69,8±2,9* 42,32±2,16* 34,8±1,75** 15,6±0,51 9,58±0,46 * 5,76±0,26*• -6 $4,6\pm0,39$ $17,4\pm1,02^*$ $16,4\pm0,92^*$ 25,1±2,46* 19,6±0,68* 11,8±0,57*• $5,3\pm0,42$ 25,3±2,54* (<0,05)

> -, , , (_ <0,05).

```
3
                                          38,5%,
                                                               66,7% ( _ <0,05).
                                                                            58,3%
            20,7% ( <0,05).
                                WOMAC
                                      , 54,2%
62,5% ( <0,01);
              14,8%
                        - 18,5%
( <sub>-</sub> <0,01).
 <0,01);
                                                                            29,6%
8,3%
                                       63%
                                            ( _<0,05).
                                 3
                                                                                -6
                                           WOMAC,
     1.
     2.
1. Hochberg M.C. Mortality in osteoarthritis. Clin. Exp. Rheumat. 2008; 26(5): 120
   -124.
             2010; 3: 35-41.
```

3.

	[]. 2012; 1. : <u>www.ukrcardio.org/journal.</u>
	php/article/670.
4.	Fernandez-Real J.M., Wifredo R. Insulin resistance and chronic cardiovascular
	inflammatory syndrome. Endocrine Rive. 2003; 24(3): 278-301.
5.	· · · · · · · · · · · · · · · · · · ·
	; 2010. 608 .
6.	Engström G., Gerhardsson de Verdier M., Rollof J. et al. C-reactive protein,
	metabolic syndrome and incidence of severe hip and knee osteoarthritis. A
	population-based cohort study. Osteoarthritis Cartilage 2009; 17(2): 168-173.
7.	,
	1 -6
	-1
	. 2009; 6: 52-56.
8.	,
	•
	2012; 2(2): 68-73.

- 9. Pasin J. S., Ferreira A. P., Saraiva A. L. et al. Diacerein decreases TNF-alpha and IL-1beta levels in peritoneal fluid and prevents Baker's yeast-induced fever in young rats. <u>Inflamm. Res.</u> 2010; 59(3): 189-196.
- 10. Bartels E.M., Bliddal H., Schøndorff P. K. et al. Symptomatic efficacy and safety of diacerein in the treatment of osteoarthritis: a meta-analysis of randomized placebo-controlled trials. Osteoarthritis Cartilage 2010; 18(3): 289-296.