

Міністерство освіти та науки, молоді та спорту України
Міністерство охорони здоров'я
Сумський державний університет
Медичний інституту



АКТУАЛЬНІ ПИТАННЯ ТЕОРЕТИЧНОЇ ТА ПРАКТИЧНОЇ МЕДИЦИНИ

Topical Issues of Clinical and Theoretical
Medicine

Збірник тез доповідей
III Міжнародної науково-практичної конференції
Студентів та молодих вчених
(Суми, 23-24 квітня 2015 року)

Суми
Сумський державний університет
2015

AEROBIC METABOLISM IN PREMATURE NEWBORNS WITH HYPOXIC-ISCHEMIC LESION OF CNS

Aloziem Nnenna Zinnah, Nwokiea Obinna- 5th year students

Scientific supervisor - associate professor Petrashenko V.O.

Sumy State University, Department of Pediatrics with Medical Genetics

All forms of oxygen deficiency are accompanied by the development of hypoxia and energetic deficiency, which leads to tension of metabolic processes of the organism. Metabolic effect of hypoxia includes stark reduce of mitochondrial activity due to a significant inhibition enzymes of the Krebs cycle: succinate dehydrogenase (SDH).

Research purpose: to increase the efficiency of diagnosis of hypoxic-ischemic CNS lesions in premature infants by determining the activity of SDG in early neonatal period.

The activity of SDH was determined in 15 conventionally healthy preterm infants (CHPI), which made the comparison group and 64 premature babies with hypoxic-ischemic CNS lesions on 1-7 day of life. They which were divided into three groups: I group – 26 premature children with mild CNS lesions; II group – 20 premature children with severe hypoxic-ischemic lesions and low birth weight; III group – 18 premature newborns with severe damage of central nervous system and extremely low birth weight.

In the premature newborn metabolic effect of hypoxia occurs in the early neonatal period through severe inhibition of mitochondrial respiratory activity, as testified by reduce activity of the main aerobic enzyme – SDH.

In infants with mild perinatal CNS injury in the early neonatal period the total number of granule of formazan into cells decreased on 33% relative to the comparison group ($p < 0,001$). In same time for infants in second and third groups were quantity 1,7 and 2,0 times lower respectively than in comparison group ($p < 0,001$).

Along with a decreased of the total number of formazan granules, were decreased number of cells containing these granules, and the average number of granules per cell.

Reduced activity of SDH indicates the initial stages of the process of decompensation, which is accompanied by disturbance of energy metabolism and leads to the formation of tissue hypoxia. At severe CNS injury further decrease activity of this enzyme shows the progression of decompensation and the formation of severe energetical disorders.

Thus, the process of aerobic glycolysis occurs in the mitochondria of brain cells with enzymes of Krebs cycle, the main of which is SDH, only in case of normal diffusion oxygen from intercellular space into neurons. So as markers of severity of perinatal hypoxic-ischemic CNS lesions in premature infants in early neonatal period can use activity of SDH.

COURSE OF ASPIRATION SYNDROME IN NEWBORN INFANTS

Chukwudifu Chibuikem Nwabueze, Mwakinpunda Leila - 6th year students

Scientific supervisor - associate professor E.K. Redko

Sumy State University, Department of Pediatrics with Medical Genetics

Aspiration syndrome of newborn children is one of reasons of development of the decompensated respiratory insufficiency that requires realization of artificial ventilation of lungs (ALV). In the department of intensive therapy new-born (ICU) of the Sumy regional child's clinical hospital for period 2009-2014 acted 1179 babies, 200 (17%) from them were hospitalized with a diagnosis neonatal aspiration. All babies were transported in the conditions of departure neonatology brigade from maternity establishments of region. During transporting ALV was conducted all children. Middle age of children at entering in the ICU was $17 \pm 0,6$ h., the 95% newborns were translated during the first twenty-four hours of life. In the ICU high-frequency ALV was conducted all children. All children by type of neonatal aspiration have been divided into two groups.

The first group included 126 infants with pure aspirating amniotic fluids (PAF), which accounted for 63% of the total number of aspiration syndromes. The second group included 74