

МІНІСТЕРСТВО ОСВІТИ ТА НАУКИ УКРАЇНИ  
СУМСЬКИЙ ДЕРЖАВНИЙ УНІВЕРСИТЕТ  
МЕДИЧНИЙ ІНСТИТУТ



**АКТУАЛЬНІ ПИТАННЯ**  
**ТЕОРЕТИЧНОЇ ТА КЛІНІЧНОЇ МЕДИЦИНИ**  
**Topical Issues of Theoretical and Clinical Medicine**

**ЗБІРНИК ТЕЗ ДОПОВІДЕЙ**  
V Міжнародної науково-практичної конференції студентів та молодих вчених  
(м. Суми, 20-21 квітня 2017 року)

Суми  
Сумський державний університет  
2017

The facies of premature infants with the signs of kidney disturbance due to severe asphyxia at 1-2 days of life can be divided into central, transitional and peripheral zones. The width of the peripheral zone ( $16,1 \pm 1,1\%$  of the radius of the drop) is almost equal to the width of the transition ( $16,7 \pm 0,9\%$  of the radius of the drop). The peripheral zone had close-grained character, and transitive - cryptocrystalline. The central part of large cross-drops contained crystals with ramifications that formed a picture "fern leaf".

**Conclusions.** Thus, the morphological picture of facies depend on the severity of asphyxia. Analysis of dried drops of urine in premature neonates with renal impairment on the background of asphyxia can be used as one of the criteria for assessing kidney function and have prognostic significance.

## QUALITY OF LIFE IN ADOLESCENTS WITH OBESITY

*Khomenko M.A.*

*Kharkiv Medical Academy of Postgraduate Education, Department of Pediatrics*

**Introduction.** Quality of life is an integral characteristic of physical, psychological and social human's functioning based on subjective perception. The urgency of studying of quality of life among patients with obesity is constantly increasing due to the growing prevalence of the disease and the influence which obesity has on the development of other chronic diseases and life expectancy.

**The purpose of the study.** Determination of quality of life in adolescents with obesity.

**Materials and methods:** A study of quality of life was conducted by questionnaire method using the 36-Item Short Form Health Survey (SF-36) in 52 adolescents with obesity (girls - 24 boys - 28) aged from 14 to 17. The questionnaire consists of 8 scales which reflect the physical and mental components of health: General Health - GH, Physical Functioning – PF, Role-Physical Functioning – RP, Bodily pain – BP, Role Emotional – RE, Social Functioning – SF, Vitality – VT, Mental Health – MH. The control group consisted of the 26 adolescents with normal body weight.

**Results:** The reduction of the physical component of health was determined in adolescents with obesity in comparison with control group: PF -  $85,4 \pm 1,4$  vs  $94,6 \pm 1,3$   $p < 0,05$ ; RP -  $77,1 \pm 1,3$  vs  $89,4 \pm 3,44$   $p < 0,05$ ; GH -  $68,6 \pm 2,5$  vs  $76,8 \pm 3,08$   $p < 0,05$ ; BP tended to decrease. A significant reduction of SF was determined among the indicators of mental components of health -  $77,8 \pm 2,8$  vs  $87,5 \pm 2,6$   $p < 0,05$ . Indicators such as VT, RE, MH tended to decrease in adolescents with obesity in comparison with control group, but did not differ significantly.

**Conclusions:** The quality of life of the adolescents with obesity was reduced mainly due to such indicators as Physical Functioning, Role-Physical Functioning, General Health, Social Functioning.

## STRUCTURE POISONING IN CHILDHOOD AND TREATMENT ALGORITHMS AT PRESENT IN SUMY CITY CHILDREN'S HOSPITAL

*Obi Okonkwo C., Antonenko I., Lito O., Goncharova A.*

*Scientific supervisor – associate prof. Romaniuk O.*

*Sumy State University, Department of Pediatrics*

**Introduction.** Acute poisoning play a significant role in the pathology of childhood.

**Aim.** Identify the structure and frequency of poisoning that occur in children, assess, depending on age, sex parents providing care of children. Characterize the features of clinical symptoms of acute poisoning, the features of laboratory diagnosis of acute poisoning in different ages of child.

**Materials and methods.** Children who were in the emergency department. Case histories of children with acute poisoning. Results of toxicology laboratory (blood, urine, vomit, food debris, etc.), morphological method results of investigation.

Work performed at the Medical Institute of Sumy State of University at the Department of Pediatrics. Under the supervision there were 234 children with acute poisoning, from 0 to 18 years, treated in Sumy City Children's Hospital for the period from 2013 to 2016. Improvement of patients