



# Wiadomości Lekarskie

**Editor in-Chief:**

Prof. Władysław Pierzchała

**Deputy Editor in-Chief:**

Prof. Aleksander Sieroń

**Statistical Editor:**

Dr Lesia Rudenko

**Managing Editor:**

Agnieszka Rosa – amarosa@wp.pl

**International Editorial Office:**

Nina Radchenko (editor)

– n.radchenko@wydawnictwo-aluna.pl

**Polish Medical Association (Polskie Towarzystwo Lekarskie):**

Prof. Waldemar Kostewicz – President PTL

Prof. Jerzy Woy-Wojciechowski – Honorary President PTL

---

**International Editorial Board – in-Chief:**

Marek Rudnicki

Chicago, USA

**International Editorial Board – Members:**

Kris Bankiewicz	San Francisco, USA	George Krol	New York, USA
Christopher Bara	Hannover, Germany	Krzysztof Łabuzek	Katowice, Poland
Krzysztof Bielecki	Warsaw, Poland	Jerzy Robert Ładny	Białystok, Poland
Zana Bumbuliene	Vilnius, Lithuania	Henryk Majchrzak	Katowice, Poland
Ryszarda Chazan	Warsaw, Poland	Ewa Małecka-Tendera	Katowice, Poland
Stanislav Czudek	Ostrava, Czech Republic	Stella Nowicki	Memphis, USA
Jacek Dubiel	Cracow, Poland	Alfred Patyk	Gottingen, Germany
Zbigniew Gasior	Katowice, Poland	Palmira Petrova	Yakutsk, Russia
Mowafaq Muhammad Ghareeb	Baghdad, Iraq	Krystyna Pierzchała	Katowice, Poland
Andrzej Gładysz	Wrocław, Poland	Tadeusz Płusa	Warsaw, Poland
Nataliya Gutorova	Kharkiv, Ukraine	Waldemar Priebe	Houston, USA
Marek Hartleb	Katowice, Poland	Maria Siemionow	Chicago, USA
Roman Jaeschke	Hamilton, Canada	Vladyslav Smiiianov	Sumy, Ukraine
Andrzej Jakubowiak	Chicago, USA	Tomasz Szczepański	Katowice, Poland
Oleksandr Katrushov	Poltava, Ukraine	Andrzej Witek	Katowice, Poland
Peter Konturek	Saalfeld, Germany	Zbigniew Wszolek	Jacksonville, USA
Jerzy Korewicki	Warsaw, Poland	Vyacheslav Zhdan	Poltava, Ukraine
Jan Kotarski	Lublin, Poland	Jan Zejda	Katowice, Poland

---

**Distribution and Subscriptions:**

Bartosz Guterman prenumerata@wydawnictwo-aluna.pl

**Graphic design / production:**

Grzegorz Sztank

www.red-studio.eu

**Publisher:**

ALUNA Publishing House

ul. Przesmyckiego 29,

05-510 Konstancin – Jeziorna

www.wydawnictwo-aluna.pl

www.wiadomoscilekarskie.pl

www.wiadlek.pl

# CONTENTS

## ORIGINAL ARTICLES

Mimoza Canga, Irene Malagnino, Alketa Qafmolla, Edit Xhajanka, Vito A. Malagnino THE IMPACT OF THE DIABETES ON ORAL HEALTH – AN OBSERVATIONAL STUDY	753
Yuliia A. Spivak, Nadiya O. Lyulka, Maksym M. Potyazhenko, Konstantin E. Vakulenko, Tetyana V. Dubrovinska BIOMARKER AND ECHOCARDIOGRAPHIC CHARACTERISTICS OF HEART FAILURE IN PATIENTS HAVING ACUTE MYOCARDIAL INFARCTION COMBINED WITH DIABETES MELLITUS OF TYPE 2	759
Maysaa Ghazi Jumaa PATTERN OF KRAS GENE EXPRESSION IN IRAQI WOMEN OVARIAN CARCINOMA	765
Petro Hasiuk, Olga Odzhubeiska, Anna Vorobets, Dmytro Korol', Tetiana Dzetsiukh, Dmytro Kindiy COMPARATIVE ANALYSIS OF ENDURANCE CEMENTS FOR THE FIXATION OF NON-REMOVABLE ORTHOPEDIC CONSTRUCTIONS UNDER THE ACTION OF CYCLIC COMPRESSION	770
Hawraa A. M. Alkhuwailidy, Muhammad M. Alrufae DNA SEQUENCING OF NOVEL YEAST ISOLATED FROM BLOODSTREAM INFECTIONS IN AL-NAJAF PROVINCE	774
Oksana S. Kapustynska, Oleg O. Samchuk, Halyna Kovalchuk, Valeriy Vdovychenko, Oleg O. Kapustynskyi, Yevgen Sklyarov, Roksolana Yaremkevych FEATURES OF COVID-19 PNEUMONIA DIAGNOSIS	781
Saly Naser Abbas, Hajer Alaa Obeid, Tahreer Shannan Alwan, Saif M. Hassan, Mahmood J. Jawad, Mohammed J. Jawad, Najah R. Hadi CORRELATION BETWEEN RS6265 SNP IN BDNF AND THE CONTEXT OF DIABETES TYPE II INVOLVEMENT IN IRAQI PATIENTS	787
Ruslana I. Falion, Yuliya I. Beketova, Yuriy O. Pospishil COMPREHENSIVE STUDY OF MANIFESTATIONS OF BRAIN TISSUE RESOLUTION IN CASE OF VARIOUS TYPES OF STROKE	791
Ahmed Zwain, Husham Qassim Mohammed EFFECT OF 20-HOUR FASTING AND LOW FAT DIET ON GHRELIN HORMONE, GLUCOSE LEVEL AND LIVER FUNCTION IN ALBINO RATS MALE	798
Mykola L. Ankin, Taras M. Petryk, Oleksander A. Radomski, Viktoria A. Ladyka, Iryna V. Kerechanyn, Larysa Y. Fedoniuk, Mykhailo P. Sas LONG-TERM RESULTS OF TREATING PATIENTS WITH OPEN FRACTURES OF LOW-LEG BONES	803
Taha Ahmed Faraj STRESS LEVELS REGARDING COVID-19 PANDEMIC AMONG NURSING STUDENTS AT UNIVERSITY OF SULAIMANI, KURDISTAN REGION, IRAQ	809
Dmytro Diachuk, Yurii Yashchenko, Iryna Zabolotna PROGNOSTIC CRITERIA OF EXCESSIVE BODY WEIGHT DEVELOPMENT AMONG SCHOOLCHILDREN BY THE RESULTS OF ANAMNESTIC SURVEY	814
Mahmood J. Jawad, Saif M. Hassan, Ahmed Kareem Obaid, Najah R. Hadi ROLE OF PRE-CESAREAN SECTION CEFOTAXIME IN AMELIORATED POST-CESAREAN COMPLICATION	818
Oleksandr V. Tsyhykalo, Nataliia B. Kuzniak, Serhij Yu. Palis, Roman R. Dmytrenko, Igor S. Makarchuk PECULIARITIES OF THE SOURCES OF ORIGIN AND MORPHOGENESIS OF THE HUMAN MANDIBLE	824
Nataliia Raksha, Oleksandr Maievskyi, Iryna Dzevulska, Rostyslav Kaminsky, Inga Samborska, Olexiy Savchuk, Oleksandr Kovalchuk PROTEOLYTIC ACTIVITY IN THE HEART OF RATS WITH HYPERHOMOCYSTEINEMIA	831
Abdulkhaleq A Ali Ghalib Al-Naqeeb, Muna A Zedian, Anaam Mohammad IMPACT OF POLYCYSTIC OVARIAN SYNDROME ON GENERAL HEALTH RELATED-QUALITY OF LIFE AMONG A SAMPLE AT "MATERNITY AND CHILDREN TEACHING HOSPITAL" IN DIWANIYAH CITY-IRAQ	836
Oleksandr Smiyan, Anastasiia Havrylenko, Andriy Loboda, Sergey Popov, Viktoriia Petrashenko, Kateryna Smiian, Tatiana Aleksakhina PECULIARITIES OF THYROID STATUS OF PRESCHOOL CHILDREN WITH ACUTE BRONCHITIS	842
Tetiana Hulai, Olena Kuzminska, Sergiy Omelchuk, Anatolii Hrynzovskyi, Tetiana Trunina, Anna V. Blagaia HYGIENIC ASSESSMENT OF THE INFLUENCE OF PESTICIDES ON THE FATTY COMPOSITION OF SUNFLOWER SEED LIPIDS	848

Marwa Jabbar, Israa Dheyaa, Khulood Abdulmahdi, Ghofran Awda NURSES' KNOWLEDGE ABOUT CORONARY ARTERY DISEASE IN AL-NASIRIYAH CITY	853
Aidyn G. Salmanov, Dmytro V. Shcheglov, Volodymyr O. Shkorbotun, Ihor M. Bortnik, Oleh Svyrydiuk, Maxim S. Gudym, Anna S. Krylova MOLECULAR EPIDEMIOLOGY OF THE TRANSMISSION OF METHICILLIN-RESISTANT <i>STAPHYLOCOCCUS AUREUS</i> IN KYIV ACUTE CARE HOSPITALS, UKRAINE	857
Olena M. Shkola, Olena V. Otravenko, Viktoriia I. Donchenko, Valeriy O. Zhamardiy, Volodymyr G. Saienko, Hanna V. Tolchieva THE INFLUENCE OF TAE-BO ON THE DEVELOPMENT OF MOTOR POTENTIAL OF STUDENTS OF MEDICAL AND PEDAGOGICAL SPECIALTIES AND ITS EFFICIENCY IN THE PROCESS OF EXTRACURRICULAR ACTIVITIES	865
Ivan M. Okhrimenko, Natalia A. Lyakhova, Valentyna V. Horoshko, Inha A. Serednytska, Svitlana S. Okhrimenko, Oleksandr L. Martenko, Svitlana V. Sprynchuk MEANS OF PSYCHOPHYSIOLOGICAL INDICATORS IMPROVEMENT OF FUTURE LAW ENFORCEMENT OFFICERS IN THE PROCESS OF THEIR SPECIALITY TRAINING	871
<b>REVIEW ARTICLES</b>	
Małgorzata Kępska-Dzilińska, Alena Zhymaila, Jolanta Malyszko KIDNEY DAMAGE IN PATIENTS AFTER ALLOGENEIC STEM CELL TRANSPLANTATION	877
Tetiana V. Fartushok, Nadiia V. Fartushok, Yu. M. Fedevych, Vladyslav V. Pyndus HISTORY OF BIOCHEMISTRY IN LVIV	881
Halyna V. Bilavych, Iryna Ja. Didukh, Viktoriia V. Stynska, Liubov M. Prokopiv, Nadiya O. Fedchyshyn, Borys P. Savchuk, Larysa Ya. Fedoniuk DEVELOPMENT OF INCLUSIVE EDUCATION IN UKRAINE IN THE CONTEXT OF WORLD TRENDS	891
<b>CASE STUDIES</b>	
Marta Yu. Mykhailevych, Oksana D. Telishevska, Ulyana D. Telishevska, Roman V. Slobodian VALUE OF ULTRASONOGRAPHY METHOD IN THE DIAGNOSIS OF TEMPOROMANDIBULAR DISORDERS AND PATIENT MANAGEMENT MONITORING. CASE REPORT	900
Anna A. Voitiuk, Tetyana A. Litovchenko, Olena N. Borodai, Nataliia A. Rudkivska DIFFERENTIAL DIAGNOSIS OF PAROXYSMAL STATES: LITERATURE REVIEW AND ANALYSIS OF A CLINICAL CASE ON THE EXAMPLE OF CLOCCS-SYNDROME IN A YOUNG MAN	907

## ORIGINAL ARTICLE

## PECULIARITIES OF THYROID STATUS OF PRESCHOOL CHILDREN WITH ACUTE BRONCHITIS

DOI: 10.36740/WLek202204117

**Oleksandr Smiyan, Anastasiia Havrylenko, Andriy Loboda, Sergey Popov, Viktoriia Petrashenko, Kateryna Smiian, Tatiana Aleksakhina**

SUMY STATE UNIVERSITY, SUMY, UKRAINE

### ABSTRACT

**The aim:** The aim of this study was to determine the characteristics of thyroid status (thyroid hormone, total and free fractions of triiodothyronine and thyroxine, reversible triiodothyronine, antibodies to thyroperoxidase) of preschool children with acute bronchitis.

**Materials and methods:** We examined 135 preschool children (from 3 to 6 years old) with acute bronchitis (main group) and 28 apparently healthy subjects who were in the control group. It used clinical-anamnestic, laboratory and instrumental research methods. Evaluation of the course of acute bronchitis was carried out in the acute period of the disease. All results were statistically processed using the SPSS 26 package.

**Results:** In 33 % of patients with acute bronchitis there are subclinical abnormalities of thyroid hormones, which manifest themselves in the form of euthyroid sick syndrome. Namely, we found an increased concentration of reversible triiodothyronine in the serum, as well as a decrease in total triiodothyronine and its free fraction.

**Conclusions:** In patients with acute bronchitis in almost every third case there are functional shifts in hormonal status, which are manifested in the form of the first variant of the euthyroid sick syndrome.

**KEY WORDS:** acute bronchitis, children, euthyroid sick syndrome, triiodothyronine, thyroxine

Wiad Lek. 2022;75(4 p1):842-847

### INTRODUCTION

An analysis of scientific publications shows that over the last century, acute infectious diseases of the respiratory system among children of all ages occupy a leading place and as a result form an important medical, social, economic and scientific problem. Respiratory tract diseases account for the vast majority, up to 90 %, of reported cases among children. According to the WHO, every year 50 % of children under the age of 5 suffer from acute respiratory infections, and from 5 to 12 years – 30 %. Acute infections of the respiratory system are 7 – 7.5 times higher than the incidence of all other infections and 1.5 – 3 times higher than in adults [1].

Acute bronchitis with its various clinical forms occupies a significant share in the structure of nonspecific acute infectious diseases of the respiratory system of children. Thus, the incidence of this nosology averages about 100 per 1,000 children [2]. Acute bronchitis is one of the five reasons for outpatient treatment by a pediatrician [3]. Among the hospitalized patients in children's infectious wards with respiratory lesions in every 2 – 4 cases in the history of the disease there is a diagnosis of acute bronchitis [4].


Acute pathological processes accompanied by inflammation, including acute bronchitis, in the course of stressful changes involve the hypothalamic-pituitary-thyroid system. Therefore, the state of thyroid status depends on the resistance and adaptive stability of the organism, which can deter-

mine the features of acute bronchitis and the consequences for the patient [5, 6, 7]. In particular, such an adaptive state as euthyroid sick syndrome (ESS) is developing, which is manifested by various combinations of changes in thyroid hormone concentrations at the subclinical level [8-10]. Most often there is the first variant of this pathology: ESS-1 or "low T3 syndrome" [8, 11]. The study of this condition will better understand the interdependence of hormonal and immune systems, their response to acute inflammatory process of the respiratory tract in children and the stress that accompanies it, as well as further use the data to predict the course of acute bronchitis and improve treatment approaches.

After analyzing the available sources of information, we found that the state of the thyroid system of preschool children with acute bronchitis remains unresolved. Therefore, we see the relevance in the study of this problem, which will improve the early diagnosis of the features of acute bronchitis and predict the consequences for the patient.

### THE AIM

The aim of this study was to determine the characteristics of thyroid status (thyroid hormone, total and free fractions of triiodothyronine and thyroxine, reversible triiodothyronine, antibodies to thyroperoxidase) of preschool children with acute bronchitis.



**Fig. 1.** Levels of thyroid hormones in the serum of the main group in the acute period of the disease

**Table I.** Condition of thyroid hormones in the serum of children with acute bronchitis (main group) in the acute phase of the disease

Hormones	N	Minimum	Maximum	Mean	
	Statistic	Statistic	Statistic	Statistic	Std. Error
T4, nmol / l	135	118,60	150,40	122,7178	0,41377
rT3, ng / dl	135	9,40	37,60	19,5733	0,68367
ft4, pmol / l	135	6,00	21,70	18,4126	0,15247
ft3, pmol / l	134	3,50	6,70	5,3597	0,10433
T3, nmol / l	135	1,01	2,74	2,0225	0,05069
TSH, mIU / l	135	1,13	3,58	1,9956	0,04281

N – number  
 Statistic – meaning  
 Minimum – minimum meaning  
 Maximum – maximum meaning  
 Mean – middle meaning  
 Std. Error – standart error

**MATERIALS AND METHODS**

We examined 135 preschool children (3 to 6 years old) with acute bronchitis (main group) who were treated in the infectious diseases departments of the Municipal Non-Profit Enterprise “Children’s Clinical Hospital of St. Zinaida” Sumy City Council and 28 healthy children (control group) who were under supervision pediatrician in the Municipal Non-Profit Enterprise “Center for Primary Health Care № 2” Sumy City Council. The first group consisted of 92 children with acute bronchitis, the second group included 43 patients with acute bronchitis with signs of ESS.

The examination of the children included in the study was agreed with their parents (informed consent) and met the requirements of bioethics, as evidenced by the protocol of the Commission on Biomedical Ethics of the Municipal Non-Profit “Children’s Clinical Hospital St. Zinaida” Sumy City Council.

Clinical and anamnestic assessments were used to comprehensively assess the patient’s thyroid status (collection of complaints, anamnesis of disease and life, physical examination), laboratory (determination by solid-phase

**Table II.** The state of thyroid hormones in children with acute bronchitis with signs of ESS and without them (main group) in the acute phase and control groups

Groups		TSH, mIU / l	T3, nmol / l	ft3, pmol / l	rT3, ng / dl	T4, nmol / l	ft4, pmol / l
I	Mean	2,0059	2,4134	6,1620	14,5478	122,6707	18,5652
	N	92	92	92	92	92	92
	Std. Error of Mean	0,04940	0,01581	0,02418	0,23458	0,52575	0,14839
II	Mean	1,9735	1,1863	3,6024	30,3256	122,8186	18,0860
	N	43	43	42	43	43	43
	Std. Error of Mean	0,08383	0,01195	0,01050	0,62251	0,65813	0,35646
Control group	Mean	1,8857	2,3732	6,2107	15,8679	121,8571	19,0357
	N	28	28	28	28	28	28
	Std. Error of Mean	0,04437	0,02619	0,03014	0,16011	0,60477	0,32060
Total	Mean	1,9767	2,0828	5,5068	18,9368	122,5699	18,5196
	N	163	163	162	163	163	163
	Std. Error of Mean	0,03637	0,04345	0,09004	0,57706	0,35834	0,13860

Mean – middle meaning

N – number

Std. Error of Mean – standart error of middle meaning

Total – total

**Table III.** Dispersion analysis of differences in the state of hormones of children with acute bronchitis with signs of ESS and without them (main group) in the acute phase and control groups

Hormones		Sum of Squares	Df	Mean Square	F	Sig.
TSH, mIU / l	Between Groups (Combined)	0,031	1	0,031	0,123	0,726
	Within Groups	33,123	133	0,249		
	Total	33,154	134			
T3, nmol / l	Between Groups (Combined)	44,124	1	44,124	2497,172	0,000
	Within Groups	2,350	133	0,018		
	Total	46,474	134			
ft3, pmol / l	Between Groups (Combined)	188,916	1	188,916	4902,456	0,000
	Within Groups	5,087	132	0,039		
	Total	194,002	133			
rT3, ng / dl	Between Groups (Combined)	7294,793	1	7294,793	835,988	0,000
	Within Groups	1160,551	133	8,726		
	Total	8455,344	134			
T4, nmol / l	Between Groups (Combined)	0,641	1	0,641	0,028	0,868
	Within Groups	3096,416	133	23,281		
	Total	3097,057	134			
ft4, pmol / l	Between Groups (Combined)	6,728	1	6,728	2,162	0,144
	Within Groups	413,820	133	3,111		
	Total	420,549	134			

Sum of Squares – sum of squares

Df – definition degrees of freedom

Mean Square – mean values of squares

F - the value of Fisher's calculation criterion

Sig. – significant - p-value (materiality)

Between Groups | (Combined) – intergroup (combined)

Within Groups – in the middle of groups (average of groups)

Total - total



enzyme-linked immunosorbent assay) of serum hormones such as thyroid-stimulating hormone (TSH), total and free (T3 and fT3) and thyroxine (T4 and fT4), reversible triiodothyronine (rT3), antibodies to thyroperoxidase), as well as instrumental research methods (chest radiography, ultrasound examination of the thyroid gland).

In children of the main group, the diagnosis of acute bronchitis was verified on the basis of complaints of children and their parents, anamnesis, objective symptoms, results of laboratory and instrumental research methods according to the clinical protocol on medical care for children in "Pediatric Pulmonology" from 13.01.2005 № 18 with changes and additions made by the order of the Ministry of Health of Ukraine from July 16, 2014 № 499 (Unified clinical protocol of primary care for adults and children "Acute respiratory infections"), Adapted clinical guidelines based on evidence, "Influenza and acute respiratory infections", 2014. Assessment of acute bronchitis, including the use of laboratory and instrumental methods, was carried out in the acute period of the disease (1 – 2 days of hospitalization).

In addition, we studied children in the main group of ESS, namely the first option, known as "low T3 syndrome". This is an adaptive state in which non-thyroid diseases with euthyroidism cause subclinical changes in serum thyroid hormone levels: decreased T3, fT3 and increased rT3.

All results were statistically processed using the SPSS 26 package. Descriptive statistics and comparisons of mean values were used to characterize the course of acute bronchitis in preschool children. Differences between groups were confirmed or refuted by analysis of variance for quantitative characteristics and the chi-square criterion, for features of nominal or rank scale.

## RESULTS

Dynamic determination of hormonal status was performed on all 135 preschool patients of the main group and 28 children of the control group. Clinically pronounced hypo- or hyperthyroidism was not detected in the examined children. With the help of ultrasound examination of the thyroid gland was ruled out the presence of organic pathology of these organs. The diagnosis of pneumonia was refuted in doubtful clinical cases after X-ray examination of the chest. Determination of the concentration of antibodies to thyroperoxidase made it possible to exclude the autoimmune etiology of changes in thyroid hormones.

The concentrations of hormones in the serum of the control group corresponded to the age norm. Their average values were: T3 –  $2.37 \pm 0.03$  nmol / l, fT3 –  $6.21 \pm 0.03$  pmol / l, T4 –  $121.86 \pm 0.60$  nmol / l, fT4 –  $19.04 \pm 0.32$  pmol / l, TSH –  $1.89 \pm 0.04$  mIU / l and rT3 –  $15.87 \pm 0.16$  ng / dl.

Compared with the indicators of the control group, the children of the main group had significant deviations and variance in the values of thyroid hormones. In general, in the acute period of the disease, the level of T3 in the serum varied from 1.01 to 2.74 nmol / l, in fT3 – from 3.5 to 6.7

pmol / l, T4 – from 118.6 to 150.4 nmol / l, fT4 – from 3.5 to 6.7 pmol / l, TSH – from 1.13 to 3.58 mIU / l and rT3 – from 9.4 to 37.6 ng / dl (table I).

Average values of concentrations of hormones of the main group of the population: T3 –  $2.02 \pm 0.05$  nmol / l, fT3 –  $5.36 \pm 0.10$  pmol / l, T4 –  $122.7 \pm 0.41$  nmol / l, fT4 –  $18.41 \pm 0.15$  pmol / l, TSH –  $1.99 \pm 0.04$  mIU / l and rT3 –  $19.57 \pm 0.68$  ng / dl (Figure 1).

Studies in children with acute bronchitis with signs of ESS and without them revealed an imbalance in changes in thyroid hormones (table II).

Analyzing the data in Tables II and III, it was found that the average level of TSH, T4 and fT4 with a probability of 95 % ( $p > 0.05$ ) did not differ statistically for the group of children with acute bronchitis with signs of ESS, without signs of ESS and in the group control.

In contrast, the level of rT3 was significantly higher in the group of children with acute bronchitis with signs of ESS ( $30.32 \pm 0.62$ ) ng / dl against ( $14.55 \pm 0.23$ ) ng / dl in the group of children with acute bronchitis without signs of ESS in the control group ( $15.87 \pm 0.16$ ) ng / dl ( $p < 0.001$ ), respectively). The level of T3 was significantly lower in the group of children with acute bronchitis with signs of ESS ( $1.18 \pm 0.01$ ) nmol / l against ( $2.41 \pm 0.02$ ) nmol / l in the group of children with acute bronchitis without signs of ESS and in the control group ( $2.37 \pm 0.03$ ) nmol / l ( $p < 0.001$ ), respectively; and fT3 in the group of children with acute bronchitis with signs of ESS ( $3.6 \pm 0.01$ ) pmol / l against ( $6.16 \pm 0.2$ ) pmol / l in the group of children with acute bronchitis without signs of ESS and in control group ( $6.21 \pm 0.01$ ) pmol / l ( $p < 0.001$ ). The above features of changes in the concentrations of thyroid hormones in the serum and gave us grounds for selection within the main group of children with manifestations of "low T3 syndrome".

The share of functional shifts in hormones is quite large. Almost every third patient ( $43 (31.85 \pm 4.02) \%$ ) showed changes in ESS.

## DISCUSSION

The problems of euthyroid sick syndrome in non-thyroid diseases are widely discussed in the modern medical literature. Thus, a number of authors observed euthyroid sick syndrome in patients with COVID-19, which was significantly associated with the severity of the disease and baseline parameters [12]. Other researchers have shown that levels of free T3 can serve as a prognostic criterion for determining the severity of the disease in the early manifestations of COVID-19 [13].

It is interesting to note that low levels of T3 are common among patients who are not in critical condition. Serum T3 levels, alone or in combination with other prognostic estimates, have been shown to be a simple and valuable tool for stratification of disease risk [14]. In addition, this hormone plays an important role in regulating the immune response, inducing protection against systemic inflammation in response to endotoxemia. Moreover, the key role of T3 is assigned to inhibit the differentiation of monocytes into macrophages [15]. At

the same time, new experimental and clinical studies show the evolutionary effect of T3 on the adaptation of affected tissue in hypoxia and on the immune response and viral load in infected tissue [16]. Some authors suggest that changes in the concentration of thyroid hormones in severe disease are a protective reaction, as it prevents excessive tissue catabolism [17].

A number of authors found that T3 was significantly correlated with lymphocyte counts in patients with bacterial sepsis, and lower hormone levels were found in patients with severe lymphopenia. In patients with severe lymphopenia, COVID-19 showed significantly lower plasma concentrations of TSH, T4, fT4 and T3 compared to patients without lymphopenia. At the same time, the indicators of inflammatory markers increased significantly: interleukin-6, C-reactive protein and ferritin [18].

Researchers believe that an increase in rT3 is a predictor of both short-term and long-term mortality. A decrease in the concentration of fT3 is a contributing factor to subsequent cardiac disorders [19].

The relationship between thyroid hormones and immune cells is complex, and T3 can model various aspects of innate and adaptive immune responses through both genomic and non-genomic mechanisms. Thyroid hormones have also been shown to affect the activity of natural killer cells and the cellular immune response [20].

Studies by Mei-Xian Xu and others (2020) have shown that children with sepsis often have euthyroid syndrome, which the authors believe is associated with high levels of interleukin-6 [21]. During the experiment, other researchers found that in patients with irritable bowel syndrome, changes in thyroid hormone levels are associated with a moderate increase in proinflammatory cytokines (tumor necrosis factor- $\alpha$ , interleukin-1 $\beta$ , interleukin-6 and interleukin-8) and decreased anti-inflammatory cytokine (interleukin-10) [22].

The problem of a clear distinction between low T3 syndrome and central hypothyroidism has been addressed by a number of researchers and proposed to determine the serological threshold of the ratio of fT3 to fT4 (pg/ml, ng/dl, respectively) at 2.0 [23].

It is possible that in the acute phase of the disease euthyroid sick syndrome is part of the body's favorable adaptation to reduce energy expenditure and activation of the innate immune response, which is important for recovery [24]. At the same time, in chronic disease, such a hypometabolic state leads to thyroid allostasis type 1, in which the amount of energy consumed and mobilized from reserves is less than the need for it [25]. To emphasize the compensatory-adaptive orientation, other researchers suggest a new name for this condition – the syndrome of adaptive thyroid imbalance. Speaking of classification, the authors point to both immobilizing (inhibitory) and mobilizing (activating) shifts in thyroid hormones [26].

Thus, the euthyroid sick syndrome occurs in a significant proportion of hospitalized patients and includes many changes in the axis of the hypothalamus-pituitary-thyroid gland. One of the hallmarks of the syndrome is a decrease in the concentration of thyroid hormones in the serum, which is often seen as an adaptive mechanism for energy savings. Cytokines released during the disease affect a number of genes involved in the metabolism of thyroid hormones, and therefore are considered the main determinants of euthyroid sick syndrome [27].

## CONCLUSIONS

Thus, in patients with acute bronchitis in the acute period of the disease in almost every third case there are functional shifts in hormonal status, which manifests itself in the form of “low T3 syndrome” (ESS-1). It should be noted that all changes are subclinical in nature, so no signs of hypo- or hyperthyroidism were detected. Thus, in children of the selected subgroup with signs of euthyroid sick syndrome, an increased concentration of reversible triiodothyronine in the serum, as well as a decrease in the total and free fraction of triiodothyronine. Our research has shown that, in response to inflammation, children with acute bronchitis may have a physiological adaptive state involving the hypothalamic-pituitary-thyroid system.

Prospects for further research are to expand the study of hormonal homeostasis, including the hypothalamus, pituitary gland and thyroid gland, to understand the role of thyroid hormones in children with acute bronchitis in age. In addition, it is important to link thyroid hormone levels to the function of the immune system and respiratory pathology. Further research in this area is needed, which is especially important in children with infectious diseases. This will allow a more accurate understanding of the pathogenetic processes of the disease and in the long run to optimize early diagnosis, the severity of acute bronchitis and predict the consequences for the patient.

## REFERENCES

1. Maidannik V.H., Falalieieva T.M., Molochek N.V., Romanenko S.Iu. Klinichni rekomendatsii z likuvannia ta profilaktyky uskladnen hostrykh respiratornykh infektsii u ditei. [Clinical guidelines for the treatment and prevention of complications of acute respiratory infections in children]. *Mizhnarodnyi zhurnal pediatrii, akusherstva ta hinekolohii*. 2019;13(3):56-99. (in Ukrainian)
2. Majdannik V.G., Emchinskaya E.A. Klinicheskie rekomendacii po diagnostike i lecheniyu ostrykh bronkhitov u detej s pozicii dokazatel'noj mediciny. [Clinical recommendations for the diagnosis and treatment of acute bronchitis in children from the standpoint of evidence-based medicine]. Kiev: NMU im. A.A. Bogomol'ca. 2014, 56p. (in Russian)
3. Lezhenko H.O., Abaturov O.Ie., Pashkova O.Ie. Rol endohennykh antymikrobynykh peptydiv u bakterialnij kolonizatsii nosohlotky v ditei iz hiperplazii retronazalnoi myhdalyny. [The role of endogenous antimicrobial peptides in bacterial colonization of the nasopharynx in children with retronasal amygdala hyperplasia]. *Zdorov'e rebenka*. 2016;6:74-81. (in Ukrainian)
4. Tokarieva N.M. Osoblyvosti mukozalnoho zakhystu pry hostrykh bronkhitakh u ditei [Features of mucosal protection in acute bronchitis in children]. [dissertation on the Internet]. Dnipro; 2019. [https://ipag-kiev.org.ua/wp-content/uploads/2019/12/dis\\_tok.pdf](https://ipag-kiev.org.ua/wp-content/uploads/2019/12/dis_tok.pdf) [date access 10.08.2021](in Ukrainian)
5. Chenchak V.A. Osobennosti dejstviya tiroksina na immunnuyu sistemu. [Features of the action of thyroxine on the immune system]. *Mezhdunarodnyj studencheskij nauchnyj vestnik* 2017;3. <https://www.eduherald.ru/ru/article/view?id=17190> [date access 10.08.2021] (in Russian)
6. Kusel'man A.I., Solov'eva I.L., Cherdancev A.P. Gerpesvirusnye infekcii u detej: rukovodstvo dlya vrachej. [Herpesvirus infections in children: a guide for doctors]. Ul'yanovsk: ULGU. 2017, 280 p. (in Russian)



7. Sirotnina O.B. Sostoyanie shchitovidnoj zhelezy u detej s timomegaliej. [Thyroid health in children with thymomegaly]. Rossijskij vestnik perinatologii i pediatrii. 2010;4:66-69. (in Russian)
8. Lee S., Farwell A.P. Euthyroid Sick Syndrome. Compr Physiol. 2016;6(2):1071.
9. Luca R.De., Davis P.J., Lin H-Y. et al. Thyroid Hormones Interaction With Immune Response, Inflammation and Non-thyroidal Illness Syndrome. Front Cell Dev Biol. 2021;8:614030. doi: 10.3389/fcell.2020.614030.
10. Ganesan K., Wadud K. Euthyroid Sick Syndrome. StatPearls Publishing. 2021, 68p.
11. Hershman J.M. Euthyroid Sick Syndrome. MSD Manual. Professional Version. <https://www.msmanuals.com/professional/endocrine-and-metabolic-disorders/thyroid-disorders/euthyroid-sick-syndrome> [date access 10.08.2021]
12. Zou R., Wu C., Zhang S. et al. Euthyroid Sick Syndrome in Patients With COVID-19. Front Endocrinol (Lausanne). 2020;11:566439. doi:10.3389/fendo.2020.566439.
13. Lui D.T.W., Lee C.H., Chow W.S. et al. Thyroid Dysfunction in Relation to Immune Profile, Disease Status, and Outcome in 191 Patients with COVID-19. J Clin Endocrinol Metab. 2021;106(2):e926-e935. doi:10.1210/clinem/dgaa813.
14. Biegelmeyer E., Scanagata I., Alves L. et al. T3 as predictor of mortality in any cause non-critically ill patients. Endocr Connect. 2021;10(8):852-860. doi:10.1530/EC-21-0080.
15. Perrotta C., Buldorini M., Assi E. et al. The Thyroid Hormone Triiodothyronine Controls Macrophage Maturation and Functions. The American Journal of Pathology. 2014;184(1):230-247. doi:10.1016/j.ajpath.2013.10.006.
16. Pantos C., Kostopanagioutou G., Armaganidis A. et al. Triiodothyronine for the treatment of critically ill patients with COVID-19 infection: A structured summary of a study protocol for a randomised controlled trial. Trials. 2020;21:573. doi:10.1186/s13063-020-04474-0.
17. Kasian V.V., Cherkun O.Iu., Sytnik D.A., Sheiko V.D. Perspektivy vyvchennia tyreoidnoho profilu u patsientiv z orhannymy dysfunktsiamy pry hostromu pankreatyti. [Prospects for the study of thyroid profile in patients with organ dysfunction in acute pancreatitis]. Visnyk problem biolohii i medytyny. 2019;1(1):43-47. doi:10.2954/2077-4214-2019-1-1-148-43-47.
18. Grondman I., de Nooijer A.H., Antonakos N. et al. The Association of TSH and Thyroid Hormones With Lymphopenia in Bacterial Sepsis and COVID-19. J Clin Endocrinol Metab. 2021;106(7):1994-2009. doi:10.1210/clinem/dgab148.
19. Razvi S., Jabbar A., Pingitore A. et al. Thyroid Hormones and Cardiovascular Function and Diseases. Journal of the American College of Cardiology. 2018;71(16):1781-1796. doi:10.1016/j.jacc.2018.02.045.
20. De Vito P., Incerpi S., Pedersen J.Z. et al. Thyroid Hormones as Modulators of Immune Activities at the Cellular Level. Thyroid. 2011;21(8):879-890. doi:10.1089/thy.2010.0429.
21. Xu M.X., Liu G., Cao L.J. et al. Zhongguo Dang Dai Er Ke Za Zhi. 2020;22(11):1215-1220. doi:10.7499/j.issn.1008-8830.2004137.
22. Babaeva A.R., Osadchuk M.A., Vidiker R.V. et al. Tireoidnaya disfunkciya i citokinovyj disbalans v patogeneze i klinike sindroma razdrzhennogo kishhechnika. [Thyroid dysfunction and cytokine imbalance in the pathogenesis and clinical manifestations of irritable bowel syndrome]. Eksperimental'naya i klinicheskaya gastroehnterologiya. 2018;(9):18-25. doi:10.31146/1682-8658-ecg-157-9-18-25. (in Russian)
23. Nomura R., Miyai K., Kuge R. et al. Free T3 to free T4 ratio less than 2.0 suggests low T3 syndrome rather than central hypothyroidism from the age of two to eighteen years. Endocr J. 2017;64(2):213-219. doi:10.1507/endocrj.EJ16-0169.
24. Langouche L., Jacobs A., Van den Berghe G. Nonthyroidal Illness Syndrome Across the Ages. J Endocr Soc. 2019;3(12):2313-2325. doi:10.1210/js.2019-00325.
25. Ruiz-Núñez B., Tarasse R., Vogelaar E.F. et al. Higher Prevalence of "Low T3 Syndrome" in Patients With Chronic Fatigue Syndrome: A Case-Control Study. Frontiers in Endocrinology. 2018;9. doi:10.3389/fendo.2018.00097.
26. Madyanov I.V., Kichigin V.A. Sindrom adaptacionnogo tireoidnogo disbalansa. Opredelenie, klassifikaciya, rasprostranennost' pri tyazhelykh somaticheskikh zabolevaniyakh. [Syndrome of Adaptive Thyroid Imbalance. Definition, Classification, Prevalence in Severe Somatic Diseases]. Bulletin of Science and Practice. 2020;6(11):217-225. doi:10.33619/2414-2948/60/26. (in Russian)
27. De Vries E.M., Fliers E., Boelen A. The molecular basis of the non-thyroidal illness syndrome. Journal of Endocrinology, 2015;225(3):R67-R81. doi:10.1530/joe-15-0133.

#### ORCID and contributionship:

Oleksandr Smiyan: 0000-0001-8225-0975 <sup>A,C,E,F</sup>  
 Anastasiia Havrylenko: 0000-0001-8237-4433 <sup>A-D</sup>  
 Andriy Loboda: 0000-0002-5400-773X <sup>A,E,F</sup>  
 Sergey Popov: 0000-0002-1789-1474 <sup>C,E,F</sup>  
 Viktoriia Petrashenko: 0000-0002-4648-8916 <sup>B,E,F</sup>  
 Kateryna Smiian: 0000-0002-8030-6418 <sup>D-F</sup>  
 Tatiana Aleksakhina: 0000-0002-9905-330X <sup>E,F</sup>

#### Conflict of interest:

The Authors declare no conflict of interest.

---

#### CORRESPONDING AUTHOR

**Oleksandr Smiyan**  
 Sumy State University  
 28 Trojtska St., 40022 Sumy, Ukraine  
 tel: +380506316005  
 e-mail: smiyana@ukr.net

**Received:** 15.12.2021

**Accepted:** 11.03.2022

---

**A** – Work concept and design, **B** – Data collection and analysis, **C** – Responsibility for statistical analysis, **D** – Writing the article, **E** – Critical review, **F** – Final approval of the article